


So smooth and natural,
everyone will notice
(but no one will know).

ASK YOUR DOCTOR
ABOUT NEW



parentheses have a place
but not on your face



Visit www.juvederm.com
or call

877-FILLER MD (877-345-5376)

NEW
FROM THE
CREATORS OF

 **BOTOX**[®]
—Cosmetic
Botulinum Toxin Type A

**BOTOX[®] Cosmetic is a
pharmaceutical drug,
available by prescription only**

Ask your doctor about new JUVÉDERM.™ To learn more, log on to www.juvederm.com or call 877-FILLER MD (877-345-5376) to find a qualified doctor in your area. **Please see important treatment considerations for patients on the following page.**

©2007 Allergan, Inc. ® mark owned by Allergan, Inc. JUVÉDERM™ mark owned by Corneal Industrie SAS. M1432-01

JUVÉDERM™ Injectable Gel

Important Treatment Considerations for Patients

In the U.S., JUVÉDERM™ injectable gel is indicated for injection into the mid-to-deep dermis for correction of moderate to severe facial wrinkles and folds (such as nasolabial folds), and is generally well tolerated. The most commonly reported side effects are temporary injection site redness, swelling, pain/tenderness, firmness, lumps/bumps, and bruising. Exposure of the treated area to excessive sun and extreme cold weather should be minimized until any initial swelling and redness have resolved.

If laser treatment, chemical peel or any other procedure based on active dermal response is considered after treatment with JUVÉDERM™ injectable gel, there is a possible risk of an inflammatory reaction at the treatment site.

Patients who are using substances that can prolong bleeding, such as aspirin or ibuprofen, as with any injection, may experience increased bruising or bleeding at injection site. You should inform your physician before treatment if you are using these types of substances. As with all skin injection procedures, there is a risk of infection. JUVÉDERM™ injectable gel should be used with caution in patients on immunosuppressive therapy, or therapy used to decrease the body's immune response, as there may be an increased risk of infection. The safety of JUVÉDERM™ injectable gel in patients with a history of excessive scarring (e.g., hypertrophic scarring and keloid formations) and pigmentation disorders has not been studied. JUVÉDERM™ injectable gel should not be used in patients who have severe allergies marked by a history of anaphylaxis or history or presence of multiple severe allergies. JUVÉDERM™ injectable gel should not be used in patients with a history of allergies to gram-positive bacterial proteins. The safety of JUVÉDERM™ injectable gel for use during pregnancy, in breastfeeding females or in patients under 18 years has not been established. The safety and effectiveness of JUVÉDERM™ injectable gel for the treatment of areas other than facial wrinkles and folds (such as lips) have not been established in controlled clinical studies.

Please go to www.juvederm.com to see a complete summary of risks and complications.